Please check the examination details below before entering your candidate information			
Candidate surname		Other names	
Pearson Edexcel International GCSE (9–1	Centre Number	Candidate Number	
Thursday 9 J	anuary	2020	
Morning (Time: 2 hours)	Paper R	eference <b>4CH1/1C 4SD0/1C</b>	
Chemistry Unit: 4CH1 Science (Double Award) 4SD0 Paper: 1C			
You must have: Calculator, ruler		Total Marks	

### Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box  $\boxtimes$ . If you change your mind about an answer, put a line through the box  $\boxtimes$  and then mark your new answer with a cross  $\boxtimes$ .

## Information

- The total mark for this paper is 110.
- The marks for each question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

### **Advice**

- Read each question carefully before you start to answer it.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P59923A
©2020 Pearson Education Ltd.



# The Periodic Table of the Elements

0 <b>He</b> helium	20 <b>Ne</b> neon 10	40 <b>Ar</b> argon 18	84 <b>Kr</b> krypton 36	131 <b>Xe</b> xenon 54	[222] <b>Rn</b> radon 86	fully
7	19 <b>F</b> fluorine 9	35.5 <b>CI</b> chlorine 17	80 <b>Br</b> bromine 35	127 	[210] <b>At</b> astatine 85	orted but not
9	16 <b>O</b> oxygen 8	32 <b>S</b> sulfur 16	79 Se selenium 34	128 <b>Te</b> tellurium 52	[209] <b>Po</b> polonium 84	ive been repo
2	14 <b>N</b> nitrogen 7	31 P phosphorus 15	75 <b>As</b> arsenic 33	122 <b>Sb</b> antimony 51	209 <b>Bi</b> bismuth 83	rs 112-116 ha authenticated
4	12 <b>C</b> carbon 6	28 <b>Si</b> silicon 14	73 <b>Ge</b> germanium 32	119 <b>Sn</b> tin	207 <b>Pb</b> lead 82	mic numbers
ဇ	11 boron 5	27 AI aluminium 13	70 <b>Ga</b> gallium 31	115 In indium 49	204 TI thallium 81	Elements with atomic numbers 112-116 have been reported but not fully authenticated
'			65 <b>Zn</b> zinc 30	112 <b>Cd</b> cadmium 48	201 <b>Hg</b> mercury 80	Elem
			63.5 <b>Cu</b> copper 29	108 <b>Ag</b> silver 47	197 <b>Au</b> gold 79	Rg roentgenium 111
			59 nickel 28	106 <b>Pd</b> palladium 46	195 <b>Pt</b> platinum 78	[271] <b>Ds</b> damstadtium 110
			59 <b>Co</b> cobalt 27	103 <b>Rh</b> rhodium 45	192 <b>                                     </b>	[268] <b>Mt</b> meitnerium 109
T hydrogen			56 Fe iron 26	101 <b>Ru</b> ruthenium 44	190 <b>Os</b> osmium 76	[277] <b>Hs</b> hassium 108
			55 Mn manganese 25	[98] <b>Tc</b> technetium 43	186 <b>Re</b> rhenium 75	[264] <b>Bh</b> bohrium 107
	nass <b>ool</b> umber		52 Cr	96 Mo molybdenum 42	184 <b>W</b> tungsten 74	[266] <b>Sg</b> seaborgium 106
Key	relative atomic mass atomic symbol name atomic (proton) number		51 <b>V</b> vanadium 23	93 <b>Nb</b> niobium 41	181 <b>Ta</b> tantalum 73	[262] <b>Db</b> dubnium 105
	relativ <b>ato</b> atomic		48 Ti titanium 22	91 <b>Zr</b> zirconium 40	178 <b>Hf</b> hafnium 72	[261] <b>Rf</b> rutherfordium 104
'			45 Sc scandium 21	89 <b>×</b> yttrium 39	139 <b>La*</b> lanthanum 57	[227] <b>Ac*</b> actinium 89
2	9 <b>Be</b> beryllium 4	24 <b>Mg</b> magnesium 12	40 <b>Ca</b> calcium 20	Sr strontium 38	137 <b>Ba</b> barium 56	[226] <b>Ra</b> radium 88
<del>-</del>	7 <b>Li</b> lithium 3	23 <b>Na</b> sodium 11	39 <b>K</b> potassium	85 <b>Rb</b> rubidium 37	133 <b>Cs</b> caesium 55	[223] <b>Fr</b> francium 87

<sup>\*</sup> The lanthanoids (atomic numbers 58-71) and the actinoids (atomic numbers 90-103) have been omitted.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.

# **Answer ALL questions.**

- 1 This question is about gases in the atmosphere.
  - (a) The box gives the names of some gases in the atmosphere.

argon cark

carbon dioxide

helium

nitrogen

oxygen

Choose gases from the box to answer these questions.

Each gas may be used once, more than once or not at all.

(i) Identify a noble gas.

(1)

(ii) Identify a gas that makes up about 78% of the atmosphere.

(1)

(iii) Identify a greenhouse gas.

(1)

(iv) Identify a gas produced by the thermal decomposition of calcium carbonate.

(1)

- (b) Sulfur reacts with oxygen to produce sulfur dioxide gas.
  - (i) Write a chemical equation for this reaction.

(1)


(ii) State an environmental problem caused when sulfur dioxide gas dissolves in water in the atmosphere.

(1)

(Total for Question 1 = 6 marks)



2 The diagram shows the electronic configuration of an atom of an element.



(a) Complete the table by giving the missing information about this atom.

(5)

name of the part of this atom labelled Z	
number of protons in this atom	
number of the group that contains this element	
number of the period that contains this element	
the charge on the ion formed from this atom	

(b) This element has three isotopes.

The table shows the mass number and percentage abundance of each isotope in a sample of this element.

Mass number	Percentage abundance (%)
24	79.2
25	10.0
26	10.8

Calculate the relative atomic mass  $(A_r)$  of this element.


Give your answer to one decimal place.


(3)

relative atomic mass = .....

(Total for Question 2 = 8 marks)

**3** The diagram shows two samples of iron, A and B.





Sample B is coated with a thin layer of zinc.

(a) Name the process used to coat iron with zinc.

(1)

(b) The two samples of iron are left outside for several weeks.

A brown solid containing hydrated iron(III) oxide forms on sample A.

(i) Give the common name for the brown solid.

(1)

(ii) Give the names of the two substances that react with the iron to form the brown solid.


(2)

1\_\_\_\_\_

2..

(c) Iron can be formed by reacting aluminium powder with iron(III) oxide.

The diagram shows how this reaction can be demonstrated.



When the magnesium fuse is lit, a very exothermic reaction occurs.

(i) State the meaning of the term **exothermic**.

(1)

(ii) The equation for the reaction between aluminium and iron(III) oxide is

$$2Al + Fe_2O_3 \rightarrow 2Fe + Al_2O_3$$

Explain what this reaction shows about the relative reactivities of aluminium and iron.

(2)

(iii) Explain why the reaction between aluminium and iron(III) oxide is a redox reaction.

(3

(Total for Question 3 = 10 marks)



- 4 This question is about ionic compounds.
  - (a) The table shows the formulae of some positive and negative ions, and the formulae of some compounds containing these ions.

	Mg <sup>2+</sup>	Al³+	NH <sub>4</sub> <sup>+</sup>
S <sup>2-</sup>	MgS	$AI_2S_3$	
NO <sub>3</sub>		Al(NO <sub>3</sub> ) <sub>3</sub>	NH <sub>4</sub> NO <sub>3</sub>
CO <sub>3</sub> <sup>2-</sup>	MgCO₃		(NH <sub>4</sub> ) <sub>2</sub> CO <sub>3</sub>

(i) Complete the table by giving the three missing formulae.

(3)

(ii) Give the name of the compound with the formula NH<sub>4</sub>NO<sub>3</sub>

(1)

(b) Sodium oxide, Na<sub>2</sub>O, is an ionic compound.

The sodium and oxide ions are held together by ionic bonds.


(i) State the meaning of the term **ionic bond**.

(2)




(ii) The diagram shows the arrangement of the electrons in a sodium atom and in an oxygen atom.

sodium atom



oxygen atom



Draw diagrams in the boxes to show the arrangement of the electrons in the ions of sodium oxide.

Include the charges on the ions.


(3)


oxide ion	sodium ion
	oxide ion

(Total for Question 4 = 9 marks)



5 The boxes show the displayed formulae of six organic compounds, P, Q, R, S, T and U.





(a) Use the letters P, Q, R, S, T and U to answer these questions.

Each letter may be used once, more than once or not at all.

- (i) Give the letter of the compound that is **not** a hydrocarbon.
  - (1)
- (ii) Give the letters of the two compounds that have the same empirical formula.
- (iii) Give the letter of the compound that is used to manufacture poly(propene).

(1)

(b) Describe a test that can be used to distinguish between compounds Q and T.	
test	
result with compound Q	
result with compound T	
(c) Compounds P, Q and R are members of the same homologous series.	
Give two characteristics of a homologous series.	(2)
1	
2	
(d) This is the displayed formula of an alkene, V.	
H H H H           C=C-C-C-H      H H	
(i) Give the name of alkene V.	(1)
(ii) Draw the displayed formula of another alkene that is an isomer of alkene V.	(1)

(e) An organic compound has the percentage composition by mass

$$H = 6.06\%$$

$$F = 57.58\%$$

(i) Show that the empirical formula of the compound is CH<sub>2</sub>F

(2)

(ii) The relative molecular mass  $(M_r)$  of the compound is 66.

Determine the molecular formula of the compound.

(2)


molecular formula = .....


(Total for Question 5 = 14 marks)

- **6** A student uses this method to investigate the reaction of dilute hydrochloric acid with zinc.
  - pour some dilute hydrochloric acid into a glass beaker
  - record the initial temperature of the acid
  - add a piece of zinc and stir the mixture
  - record the temperature of the mixture after one minute
  - (a) Write a word equation for the reaction of dilute hydrochloric acid with zinc.

(1)

(b) The diagram shows the thermometer readings for this reaction.





Complete the table, giving all values to the nearest 0.1°C.

(3)

temperature in °C after adding zinc	
temperature in °C before adding zinc	
temperature change in °C	

(c)	Another student repeats the method using five different metals to compare their r	eactivity.
	(i) This student uses a polystyrene cup instead of a glass beaker.	
	Explain why a polystyrene cup is better than a glass beaker in this investigation	n. (2)
	(ii) Give three factors that the student should keep constant in this investigation.	(3)
1		
2		
3		

(d) The table shows some of the student's results.

Metal added	Observation	Temperature change in °C
copper	no bubbling	0.0
iron	slow bubbling	
magnesium	rapid bubbling	8.7
tin	very slow bubbling	1.4
zinc	moderate bubbling	5.1

(1)

(ii) Predict the temperature change for iron.

(1)

temperature change = ......°

(iii) Deduce the order of reactivity of the five metals.

(1)

most reactive .....

.....

.....

least reactive

(Total for Question 6 = 12 marks)



- 7 This question is about some of the halogens and their compounds.
  - (a) (i) Which element is a liquid at room temperature?

(1)

- A astatine
- **B** bromine
- C chlorine
- **D** iodine
- (ii) Which element has the palest colour?

(1)

- A astatine
- **B** bromine
- C chlorine
- **D** iodine
- (iii) Which element is the least reactive?

(1)

- A astatine
- **B** bromine
- **C** chlorine
- **D** iodine

(b) A teacher uses displacement reactions to demonstrate the reactivities of some halogens.

She adds solutions of chlorine, bromine and iodine separately to three different sodium halide solutions.

The table shows some of the teacher's results.

	sodium chloride	sodium bromide	sodium iodide
chlorine solution	not done	solution turns orange	
bromine solution	solution stays orange	not done	solution turns brown
iodine solution		solution stays brown	not done

A change in colour of the solution indicates that a reaction has occurred.

(i) Complete the table by predicting the missing results.

(2)

(ii) State why the teacher does not add bromine solution to sodium bromide solution.

(1)

(iii) The word equation for the reaction of bromine with sodium iodide is  $bromine \ + \ sodium \ iodide \ \rightarrow \ iodine \ + \ sodium \ bromide$ 

Write a chemical equation for this reaction.

(1)



(c) A technician sees an unlabelled bottle conta	ining a liquid.
He knows that the liquid is a solution of one	of these compounds.
<ul><li>copper(II) chloride</li></ul>	
<ul> <li>copper(II) bromide</li> </ul>	
<ul><li>iron(II) chloride</li></ul>	
<ul><li>iron(II) bromide</li></ul>	
Describe chemical tests that the technician	could use to identify the compound in
the solution.	(6)
	(Total for Question 7 = 13 marks)

**BLANK PAGE** 




8	<ul><li>(a) Carbon dioxide changes directly from a solid to a gas without becoming a liquid.</li><li>(i) Give the name of the change of state from solid to gas.</li></ul>	(1)
	(ii) Describe the test for carbon dioxide gas.	(2)
	(b) Carbon dioxide is a simple molecular covalent substance.	
	(b) Carbon dioxide is a simple molecular covalent substance.  Explain why carbon dioxide turns from a solid to a gas at a very low temperature.	(2)
		(2)
		(2)


(c) Diamond and graphite are both giant covalent substances made	le up of carbon atoms.
<ul> <li>diamonds are used in cutting tools</li> </ul>	
<ul> <li>graphite is used in pencils to make marks on paper</li> </ul>	
Explain, with reference to structure and bonding, why each sub	ostance is suitable
for its particular use.	
	(6)
(Total for Q	uestion 8 = 11 marks)



**BLANK PAGE** 







This is the student's method.

- measure the mass of the spirit burner and fuel
- add 100 cm<sup>3</sup> of water to the copper can
- record the temperature of the water
- use the spirit burner to heat the water until the temperature rises by 30 °C
- immediately measure the new mass of the spirit burner and fuel

(a)	Suggest why the student measures the mass of the spirit burner and fuel immed	iately
	after heating the water.	

(1)

- (b) When the fuel is burned, the student notices that a black solid forms on the bottom of the copper can.
  - (i) Identify the black solid.

(1)

(ii) Explain why the black solid forms.

(2)



(c) (i) Show that the heat energy change, Q, to raise the temperature of 100 cm<sup>3</sup> of water by 30 °C is approximately 13 kJ.

[mass of 
$$1.0 \,\mathrm{cm}^3$$
 of water =  $1.0 \,\mathrm{g}$ ]

[c for water = 
$$4.2 \text{ J/g/}^{\circ}$$
C]

(3)

(ii) The student burns  $0.96\,\mathrm{g}$  of methanol,  $CH_3OH$ 

Calculate the molar enthalpy change,  $\Delta H$ , in kJ/mol, for the combustion of methanol.

Include a sign in your answer.

$$[M_r \text{ of methanol} = 32]$$

(3)

 $\Delta H = \dots kJ/mol$ 

(d) The table shows data book values for the molar enthalpy change,  $\Delta H$ , for the combustion of some alcohols with different numbers of carbon atoms per molecule.

Number of carbon atoms per molecule	1	2	3	4	5
Molar enthalpy change, $\Delta H$ , in kJ/mol	-730	-1370	-2020	-2680	-3320




(2)

(i) Plot the data values from the table on the grid.

Draw a straight line of best fit.

Number of carbon atoms per molecule



(ii) Deduce the value of  $\Delta H$  for an alcohol with six carbon atoms per molecule.

Show on the graph how you obtained your answer.

(2)

 $\Delta H = k I/mc$ 

(iii) State the relationship between  $\Delta H$  and the number of carbon atoms per molecule.

(1)

(Total for Question 9 = 15 marks)

10	Nitric acid	(HNO₃) is used in the production of fertilisers.		
	Nitric acid is manufactured in three stages.			
	Stage 1	ammonia reacts with oxygen in the presence of a platinum catalyst to produce nitrogen monoxide gas, NO, and water.		
	Stage 2	nitrogen monoxide gas reacts with more oxygen to produce nitrogen dioxide gas, $NO_2$ .		
	Stage 3	nitrogen dioxide gas reacts with water to produce nitric acid and more nitrogen monoxide gas.		
	(a) (i) Coi	mplete the chemical equation for the reaction in stage 1.	(1)	
		$NH_3 + \dots NO + \dots H_2O$		
	(ii) Giv	re the meaning of the symbol $\rightleftharpoons$	(1)	
	(iii) Sta	te the purpose of the platinum catalyst.	(1)	
	(b) Give a	chemical equation for the reaction of nitrogen monoxide and oxygen in sta	age 2.	

(c) (i) The equation for the reaction in stage 3 is

$$3NO_2 + H_2O \rightarrow 2HNO_3 + NO$$

Calculate the maximum mass, in tonnes, of nitric acid that could be produced in this reaction from 11.5 tonnes of nitrogen dioxide.

$$[1 \text{ tonne} = 1.0 \times 10^6 \text{g}]$$

(4)

mass of nitric acid = .....tonnes

(ii) Suggest what use can be made of the nitrogen monoxide gas formed in stage 3.

(1)

**QUESTION 10 CONTINUES ON NEXT PAGE** 

(d) When copper(II) oxide reacts with dilute nitric acid, copper(II) nitrate is produced.

The equation for the reaction is

$$CuO + 2HNO_3 \rightarrow Cu(NO_3)_2 + H_2O$$

0.200 mol of nitric acid reacts with excess copper(II) oxide.

A mass of 15.3 g of copper(II) nitrate is produced.

Calculate the percentage yield of copper(II) nitrate.

 $[M_r \text{ of copper(II) nitrate} = 187.5]$ 

(3)

percentage yield = ..... %

(Total for Question 10 = 12 marks)

**TOTAL FOR PAPER = 110 MARKS**